

Assessment of mother's knowledge about neonatal danger sign and associated factors in Sodo town, Wolaita Zone, Southern Ethiopia

Molalegn Mesele Gesese^{1*},
Yohannes Fikadu Geda²

¹Department of Midwifery, College of health science and medicine, Wolaita Sodo University, Wolaita Sodo, Ethiopia

²Department of Midwifery, College of Medicine and Health Science, Wolkite University, Wolkite, Ethiopia

Abstract

Background: In Ethiopia neonatal mortality rate was 30 deaths per 1,000 live births in 2019. The survival of the child is determined at neonatal period. Neonatal danger sign refers to the occurrence of sign which show newborn mortality and morbidity. To reduce mortality and morbidity initial therapeutic intervention is necessary. Thus, family should be recognized neonatal danger sign and bring the newborn infant to health facility.

Objective: the aim of the study was to assess knowledge about neonatal danger signs and associated factors among mothers in Sodo town, Wolaita Zone, southern Ethiopia.

Method: community based cross-sectional study design was employed from October 1st to 30th, 2019. Systematic sampling technique was used to select 410 mothers. A pre-tested, structured and interviewed-administered questionnaire was used to collect data. Data were entered using Epi-data version 3.1 and analyzed using SPSS version 21. Bivariate and multivariable analysis was carried out using binary logistic regression to test and check the association between dependent and independent variables.

Result: From 410 participants, 275(67.1%) (95% CI: 66.2%, 69.1%) of mothers had good knowledge regarding neonatal danger signs. 226(54.9%) of mothers identified diarrhea as neonatal danger signs. Husband's educational status secondary school (AOR: 3.84, 95% CI: 1.66,8.89) and college and above (AOR: 2.41, 95% CI: 1.16, 5.03), place of residence(AOR: 2.04, 95% CI: 1.16, 3.57), Antenatal care service utilization (AOR: 2.79, 95 % CI: 1.49,5.19), Postnatal care service utilization (AOR: 1.87, 95% CI: 1.18,2.94) and mothers heard about neonatal danger signs (AOR: 2.14, 95% CI: 1.29,3.54) were factors significantly associated with mothers knowledge about neonatal danger signs.

Conclusion and recommendation: Although not satisfactory in views of expectation, relatively high level of mothers' knowledge about neonatal danger sign neonatal danger signs had been observed in study area as compared with previous reports. The government, town health office, health workers should contribute to create awareness about neonatal danger signs in the community.

Keywords: Ethiopia, knowledge, neonatal danger sign, Sodo town

***Corresponding author:** Molalegn Mesele, Department of Midwifery, College of medicine and health science, Wolaita Sodo University, Wolayita Sodo, Ethiopia. Address: Tel: +251-932214732 E-mail: emimolalegn@gmail.com.

Citation: Molalegn MG, Yohannes FG (2021) Assessment of mother's knowledge about neonatal danger sign and associated factors in Sodo town, Wolaita Zone, Southern Ethiopia. J Prev med Vol. 6 No. 3: 80

Received: February 26, 2021; **Accepted date:** February 05, 2021; **Published date:** March 12, 2021

Introduction

In human life, the first 28 day of life is known as neonatal period. At this stage the child is at highest risk of dying. The survival of the child is determined at neonatal period. It is crucial that appropriate feeding care are providing during this period. It is very critical time to improve the child's chances of survival and to lay the foundations for a healthy life. In order to continue

to accelerate the reduction in under-five mortality, focusing on newborns should be a primary concern [1, 2].

Between 1990 and 2017 the global Neonatal Mortality Rate (NMR), is decreased by 51% from 36.6 deaths per 1,000 live births in 1990, to 18.0 deaths per 1,000 live births in 2017. The estimated numbers of neonatal death during the same period decreased from 5.0 million to 2.5 million. Annual NMRs vary widely across the world, but west and central Africa and south

Asia had the highest NMRs in 2017 [3]. Sub-Saharan Africa had the highest NMR in 2018 at death per 1,000 live births; which is 10 times more likely to die than a child born in a high income country [4]. In Ethiopia, neonatal mortality rate was 30 deaths per 1,000 live births in 2019; in 2016, it was 29 [5, 6].

Neonatal danger sign refers to the occurrence of sign which show great danger of newborn mortality and morbidity and necessity for initial therapeutic intervention. Convulsion, which is one of newborn danger sign occur because of unexpected, unusual electrical activity in the brain. Fever is a raise of body temperature above the normal regular variation of the famous manifestation of diseases and it is the most common causes to find health care provider and visit physician in childhood. Lethargy and poor breast milk sucking, especially in a baby who was feeding well previous, are very imperative and sensitive indicators of neonatal sickness. An increased respiratory rate (more than sixty per minute while counted for a minimum one minute) and chest retractions indicate a serious problem. Jaundice and Vomiting are also very significant danger signs which require urgent treatment [7-10].

In Ethiopia childhood mortality levels are decreasing. Ethiopian Demographic and health survey 2019 indicate that; neonatal mortality rate are 30 deaths per 1,000 live births. Post neonatal mortality rate was 13 deaths per 1,000 live births. Infant mortality rate was 43 deaths per 1,000 live births and child mortality rate was 12 deaths per 1,000 children surviving to age 12 month [5].

Ethiopia has made an important work on health care implementation through integrated child care program with health extension workers. But, still pain from mortalities and morbidities related with neonatal danger signs. This is mostly related with lack of knowledge about neonatal danger signs [11-13]. The health extension program is highly essential in delivering quality maternal, neonatal and child health services through efficient and effective linkages between health center, health post and community. Solitary limited studies remained conducted in Ethiopia with respect to care seeking practice of mothers for their children. So this research intended to assess the mothers' knowledge about neonatal danger signs.

Materials and Methods

Study setting and design

Community based cross-sectional study was conducted on mothers who give birth within 12 months in Sodo town, Wolaita Zone, Southern Ethiopia, from October 1st to 30th, 2019. Sodo town is located 390 km south of Addis Ababa, the capital city of Ethiopia and 153 kilometer far from Regional city, Hawasa. The town is divided in to four administrative sub-cities. As the town administrative office report, the total population of the town in 2018 was 182,607 (93,130 male and 89,477 females) from these, 28,499 under-five age children and 4576 infants less than one year of age. There are also 4963 women in the reproductive age group (15-49 years). Functioning health facilities in the town includes 17 health posts, 17 medium and lower level clinics, 3 health centers

and two Hospitals (one private and one government).

Source and study population

All women in reproductive age group (15-49) living in Sodo town and two Sodo zurea kebeles were the population of study. Among them, lists of mothers who give birth in the last 12 months were selected. Preceding the survey was obtained from the selected community with the help of health extension workers.

Sample size determination and sampling procedure

Sample size was calculated by using the single population proportion formula by taking the following consideration; marginal error of 0.05, with 95% confidence interval and p-value 41% from study conducted in Arba Minch General Hospital, Southern, Ethiopia cross-sectional study in title knowledge about neonatal danger sign and associated factors among mothers attending immunization clinic in Arba Minch general hospital [14]. By adding non response rate 10%, the final sample size was 410.

Mothers who give birth in the last 12 months was coded with the help of health extension workers and the sample size was allocated proportionally to all administrative sub-cities of Sodo town and two Sodo zurea kebeles, then by using systematic random sampling technique every 6th mothers were interviewed.

Measurement

Data was collected by face to face interview method with structured questionnaire. The questionnaire was first developed in English and translated into Amharic. The questionnaire items include; socioeconomic and demographic characteristics, obstetric and knowledge on neonatal danger sign. The questionnaire was taken from maternal and child health program of Johns Hopkins Program for International Education in Gynecology and Obstetrics. The outcome of this study is mothers' knowledge about neonatal danger signs [15].

Ethics approval and consent to participate

Ethical approval was obtained from Wolaita Sodo University College of health science and medicine Ethical review board. Permission was obtained from Sodo town health office.

Consent to participate: A written informed consent was obtained from the parents of the minors and study participants. The purpose and objective of the study was well explained to study participants and they were informed about their full right to withdrawal or discontinue participation at any time they want. In addition privacy and confidentiality of participant was kept.

Result

Socio-demographic characteristics

From a total of 412 mothers selected to participate in this study 410 (100% response rate) were completed the interview. The mean age was 24.45 (SD \pm 6.2) years. Majority of mothers

173(42.2%) have primary (grade 1 -8) educational level. 295(72%) mothers were house wife).

Obstetrics history of participants

Regarding obstetric history of respondents; three hundred fifty four (86.3%) of them attend Antenatal Care (ANC) follow up for their last pregnancy. 223(54.4%) have history of Postnatal Care (PNC) services utilization.

Mother's knowledge about neonatal danger signs

The overall knowledge level of mother on neonatal danger sign indicated that 275(67.1%) (95% CI: 66.2%, 69.1%) have a good knowledge regarding neonatal danger signs.

From a total of 410 participants, majority of mothers 284(69.3%) had information or heard about neonatal danger sign. Among listed neonatal danger sign, high awareness among mothers was, diarrhea 226(54.9%), persistent vomiting 211(51.2%) and followed by fever 210(51.0%).

Factors associated with maternal knowledge on neonatal danger signs

In multivariable logistic regression, husband educational status, residence, ANC follow up, PNC follow up and heard about neonatal danger signs were factors contributed for mothers knowledge with neonatal danger sign. Mothers whose husband educational status from 9 to 12(secondary education) has 3.84 time (AOR: 3.84, 95% CI: 1.66, 8.89) and college and university 2.41 time (AOR: 2.41, 95% CI: 1.16, 5.03) more likely have good knowledge level than when compared with never attend school. Mothers who were lived in urban were 2.04 time more likely knowledgeable than when compared with mothers who lived in rural (AOR: 2.04, 95% CI: 1.16, 3.57).

Discussion

Reducing child illness and death required abrupt mothers and care givers recognition of suggestive known danger signs to take immediate actions. The finding this study indicates that the level of good maternal knowledge about neonatal danger sign in Sodo town was 275(67.1%) (95% CI: 66.2%, 69.1%). The prevalence of mother's good level of knowledge about neonatal danger sign was higher when compare with the study conducted in Arba Minch general hospital, Arba Minch town 41%(14), study done in Wolkite town, Gurage zone, southern Ethiopia 31.3%(18), study done in Chench district, southern Ethiopia 50.3% and study done in Nigeria 30.3%. The difference might be due to time gap, sample size variation, socio-cultural and residence variation.

The current study revealed that husband's educational level was significantly associated with knowledge of mothers towards neonatal danger sign. Husband education was significant predictor of mothers' good knowledge about neonatal danger signs were 3.84 times among mothers whose husbands achieved secondary school (grade 9 to 12) and 2.41 times among mothers whose husbands achieved college and university educational

level. This study was in line with a study done in North West Ethiopia [16]. This might be due to that as fathers' in culturally pre-dominant position in decision-making positively affected the knowledge and attitude of mothers. Place of residence was significantly associated with level of mothers knowledge about neonatal danger sign. Mothers lived in urban area was 2.04 time more likely had good knowledge of neonatal danger sign than mothers lived in rural area. The current study is in line with a study done in Chench district. The reason might be mothers who live in urban had more access to health information from different sources and more to seek health care as compare to mother who live in rural part.

The current study also indicated that ANC follow up creates a good opportunity for mother to had good knowledge towards neonatal danger sign. Mothers attended ANC follow up were 2.79 times more likely to had good knowledge about neonatal danger sign as compared to their counterparts. The finding of this study is consistent with a study conducted in North West Ethiopia [16]. The reason might be during ANC the health care provider gives information and counseled on neonatal danger sign this increase mother's knowledge about neonatal danger sign. PNC follow up where statistically significant factor associated with mothers' knowledge on neonatal danger sign. Mothers had PNC follow up had 1.87 times more likely to have good knowledge when compared with to those mothers who had no PNC follow up. This study was in line with a study done in Wolkite town, Gurage Zone, Southern Ethiopia, North West Ethiopia and Arba Minch general hospital [14, 16, 18]. The possible reason might be mothers had PNC follow up are counseled about neonatal danger sign; this increase knowledge of mothers' related neonatal danger signs.

This study indicated that, mothers heard about neonatal danger signs were factor significantly associated with mothers' knowledge about neonatal danger signs. Those mothers had heard about neonatal danger signs have 2.14 times more likely to had good knowledge about neonatal danger signs. The reason may be the awareness creation or those mothers heard about neonatal danger signs are knowledgeable about neonatal danger signs.

The limitation of this study was cross-sectional nature of study design which leads the assessments of the exposure and outcome at the same point in time, so that we cannot formulate a cause and effect relationship between identified factors and mothers knowledge about neonatal danger signs.

Conclusion

Although not satisfactory in views of expectation, relatively high level of mothers' knowledge about neonatal danger sign had been observed in study area as compared with previous reports. Mothers' husband educational status, ANC and PNC follow up, place of residence and heard about neonatal danger signs are factors associated with neonatal danger sign.

The current study suggested that as part of health education and sensitization, women should be taken through counseling on

danger signs of newborn ill health preceding to their discharge from hospital so that they can easily detect signs and hurry to health care facilities whenever necessary. The government, town health office, health workers and Nongovernmental organization should contribute to create awareness about neonatal danger signs in the community.

Authors' contributions

MMG: wrote the proposal, participates in data collection, analysis the data, made revision on proposal and wrote the manuscript. YFG: made revision on manuscript and participate on data analysis.

Acknowledgment

We would like to express our heartfelt thanks for Wolaita Sodo University, all individual for their contribution and Wolaita Sodo health office for their cooperation.

References

- Graves, MF (2016) The vocabulary book: Learning and instruction. Teachers College Press.
- UNICEF (2014) Committing to child survival: A promise renewed.
- Hug L, Alexander M, You D, Alkema L (2019) National, regional and global levels and trends in neonatal mortality between 1990 and 2017, with scenario-based projections to 2030: A systematic analysis. *Lancet Glob Health* 7:10-20.
- WHO (2018) State of global health, newborns: Reducing mortality.
- Ethiopian, PublicHealthInstitute (2019) Ethiopian Mini Demographic and Health Survey 11.
- Senarath U, Fernando DN, Rodrigo I (2007) Effect of training for care providers on practice of essential newborn care in hospitals in Sri Lanka. *J Obstet Gynecol Neonat Nurs* 36:531-541.
- Hadush A, Kassa M, Kidanu K, Tilahun W (2016) Assessment of knowledge and practice of neonatal care among post natal mothers attending in Ayder and Mekelle Hospital in Mekelle, Tigray, Ethiopia 2013. *SMU Med J* 3:509-525.
- Baqui AH, Rahman M, Zaman K, El Arifeen S, Chowdhury HR, et al. (2007) A population-based study of hospital admission incidence rate and bacterial aetiology of acute lower respiratory infections in children aged less than five years in Bangladesh. *J Health Popul Nutr* 25:179.
- Young Infants Clinical Signs Study Group (2008) Clinical signs that predict severe illness in children under age 2 months: a multicentre study. *The Lancet* 371:135-142.
- Assefa T, Belachew T, Tegegn A, Deribew A (2008) Mothers' health care seeking behavior for childhood illnesses in derra district, northshoa zone, oromia regional state, ethiopia. *Ethiopian J Health Sci* 18.
- Devi SK (2015) A Study to Assess the Effectiveness of Video Assisted Teaching Module (VATM) on Knowledge Regarding Care of Newborn Baby under Photo Therapy among Female Health Workers in Selected Hospital, Bhubaneswar, Odisha. *Int J Adv Nurs Sci Prac* 2:29-35.
- Dongre AR, Deshmukh PR, Garg BS (2009) Awareness and health care seeking for newborn danger signs among mothers in peri-urban Wardha. *Ind J Pediatr* 76:691-693.
- Degefa N, Diriba K, Girma T, Kebede A, Senbeto A, et al. (2019) Knowledge about Neonatal Danger Signs and Associated Factors among Mothers Attending Immunization Clinic at Arba Minch General Hospital, Southern Ethiopia: A Cross-Sectional Study. *Hindawi Bio Med Res Int* 3.
- Maternal JH (2001) Neonatal health (MNH) Program. Birth preparedness and complication readiness: A matrix of shared responsibilities. *JHPIEGO-MNH* 23-31.
- Nigatu SG, Worku AG, Dadi AF (2015) Level of mother's knowledge about neonatal danger signs and associated factors in North West of Ethiopia: A community based study. *BMC research notes* 8:1-6.
- WHO (2012) Caring for the newborn at home. A training course for community health workers; community health workers manual, Geneva.
- Anmut W, Fekecha B, Demeke T (2017) Mother's knowledge and Practice About Neonatal Danger Signs and Associated Factors in Wolkite Town, Gurage Zone, SNNPR, Ethiopia. *J Biomed Sci* 6:4-33.